

## Case Report

# Chest wall reconstruction using a combined musculocutaneous anterolateral–anteromedial thigh flap

Pearlie W. W Tan, Chin-Ho Wong, Heng-Nung Koong<sup>1</sup>, Bien-Keem Tan

Department of Plastic, Reconstructive and Aesthetic Surgery, <sup>1</sup>Department of General Surgery, Singapore General Hospital, Singapore.

**Address for correspondence:** Bien-Keem Tan, FRCS (Plast). Singapore General Hospital, Outram Road, Singapore - 169 608.  
E-mail: bienkeem@singnet.com.sg

### ABSTRACT

We present a massive 25 cm x 20 cm chest wall defect resulting from resection of recurrent cystosarcoma phyllodes of the breast along with six ribs exposing pleura. The chest wall was reconstructed with a Prolene mesh–methylmethacrylate cement sandwich while soft tissue reconstruction was carried out using a combined free anterolateral–anteromedial thigh musculocutaneous flap with two separate pedicles, anastomosed to the thoracodorsal and thoracoacromial vessels, respectively. We explain our rationale for and the advantages of combining the musculocutaneous anterolateral thigh with the anteromedial-rectus femoris thigh.

### KEY WORDS

Anterolateral thigh flap; Anteromedial thigh flap; Chest wall defects; Combined free flap

### INTRODUCTION

For a large chest wall defect measuring 25 cm x 20 cm, resulting from the excision of a recurrent Cystosarcoma Phylloides and involving ribs 4 to 9 with underlying pleura, the skeletal stabilization was provided by a proline mesh - methymethacrylate sandwich and the soft tissue repair was done by a combined Antero-lateral and Antero-medial thigh flap on two separate vascular pedicles. Along with the case description we are attempting to present the vascular anatomy of this dual flap.

### CASE REPORT

A 52-year-old woman presented with recurrent cystosarcoma phyllodes of the left breast 12 years after primary resection [Figure 1].

Intra-operatively, a large tumour measuring 12 cm transversely was found adherent to the chest wall, invading the skin, ribs and pleura. The thoracic surgeon resected the entire left anterolateral chest wall from the



Figure 1: Recurrent Cystosarcoma Phyllodes of the anterolateral chest wall




Figure 2: (A) Prolene mesh–Methylmethacrylate sandwich prosthesis. (B) Flap inset and revascularisation




Figure 3: (A) Lateral circumflex femoral system supplying the Rectus Femoris and Vastus Lateralis muscles and (B) Cadaveric specimen showing the design of the flap raised. Note the pedicle to the Anteromedial Thigh (Yellow Pointer) arising from the descending branch

midsternum to the midaxillary line, comprising skin, intercostal muscles and ribs 4–9 with underlying pleura, leaving a 25 cm x 20 cm defect. Both internal mammary arteries were ligated during resection.

Skeletal reconstruction was performed with a Prolene mesh–methymethacrylate cement sandwich [Figure 2], followed by immediate soft-tissue reconstruction. We harvested a combined musculocutaneous anterolateral thigh (ALT) with anteromedial thigh (AMT) flap, including

the vastus lateralis and rectus femoris muscles, giving us two separate pedicles.

The flap was revascularised using two sets of recipient vessels, namely the thoracodorsal and the thoracoacromial arteries. Extra venous drainage was facilitated by a long saphenous vein graft [Figure 2]. The donor site was skin grafted.

Postoperatively [Figure 3], the patient underwent



Figure 4: Six months postoperatively

radiotherapy. She was able to ambulate with a walking aid and lived for another 1 year before she succumbed to distant metastases.

## DISCUSSION

The options for large chest wall defects range from local, pedicled (latissimus dorsi, pectoralis major, external oblique or rectus abdominis) muscle or free tissue transfer (single or multiple) according to Chang *et al.*<sup>[1]</sup> In our case, the ipsilateral latissimus dorsi could not be used as it was transected previously. The TRAM flap was precluded as both internal mammary vessels were ligated, and chasing what was left as a recipient site for a free TRAM would be challenging. Abdominal closure of a large TRAM would also lead to tension on the defect. Other free flap options were a large thigh flap or a contralateral latissimus dorsi flap. The latter was not chosen as it would require repositioning the patient while surgery.

Anterior thigh flaps provide bulk for contour deficiencies and excellent coverage of implants.<sup>[2]</sup> The ALT flap has been popularized for its versatility and its vascular anatomy is well described.<sup>[3-6]</sup> Most surgeons would limit its dimensions to 24 cm x 12 cm. To increase its dimension, we combined it with an AMT flap, resulting in a 25 cm x 20 cm skin paddle. The AMT flap is supplied by a branch of the descending branch of the lateral circumflex femoral artery (described by 73% of clinical series), whereas other vascular variations are perforators arising from the lateral circumflex femoral artery,<sup>[7]</sup> the superficial femoral artery<sup>[8,9]</sup> and from minor muscle branches to the rectus femoris.<sup>[10]</sup> Our flaps were derived from the descending branch [Figure 4], but we separated it into two pedicles to improve circulation

and safeguard against thrombosis. We included the vastus lateralis and rectus femoris to maximise muscle coverage, to improve local vascularity and offset the hard palpability of the underlying cement prosthesis. In addition, the tendinous portions of these muscles were anchored to the mesh–methymethacrylate sandwich to prevent the flap from being dragged down by gravity. Lastly, we chose not to raise it as a perforator flap as experience has shown, that very large ALT flaps, based on even two to three perforators can still show marginal necrosis and, more importantly, laying bare perforators against a cement base would risk vessel spasm.

This patient had successful coverage of a large cement prosthesis with preservation of thoracic function. The main drawback was knee weakness from sacrifice of the vastus lateralis and rectus femoris muscles.

## ACKNOWLEDGMENT

We thank Dr. Chew Khong Yik for helping us with the dissection of the cadaveric specimen.

## REFERENCES

1. Chang RR, Mehrara BJ, Hu QY, Disa JJ, Cordeiro PG. Reconstruction of complex oncologic chest wall defects: A 10-year experience. *Ann Plast Surg* 2004;52:471-9.
2. Ao M, Uno K, Maeta M, Nakagawa F, Saito R, Nagase Y. De-epithelialised anterior (anterolateral and anteromedial) thigh flaps for dead space filling and contour correction in head and neck reconstruction. *Br J Plast Surg* 1999;52:261-7.
3. Song YG, Chen GZ, Song YL. The free thigh flap: A new free flap concept based on the septocutaneous artery. *Br J Plast Surg* 1984;37:149-59.
4. Wong CH, Wei FC, Fu B, Chen YA, Lin JY. Alternative vascular pedicle of the anterolateral thigh flap: The oblique branch of the lateral circumflex femoral artery. *Plast Reconstr Surg* 2009;123:571-7.
5. Koshima I, Yamamoto H, Hosoda M, Moriguchi T, Orita Y,

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50

Nagayama H. Free combined composite flaps using the lateral circumflex femoral system for repair of massive defects of the head and neck regions: An introduction to the chimeric flap principle. *Plast Reconstr Surg* 1993;92:411-20.

6. Baek SM. Two new cutaneous free flaps: The medial and lateral thigh flaps. *Plast Reconstr Surg* 1983;71:354-65.

7. Koshima I, Soeda S, Yamasaki M, Kyou J. The free or pedicled anteromedial thigh flap. *Ann Plast Surg* 1988;21:480-5.

8. Koshima I, Hosoda M, Inagawa K, Moriguchi T, Orita Y. Free medial thigh perforator-based flaps: New definition of the pedicle vessels and versatile application. *Ann Plast Surg* 1996;37:507-15.

9. Xu Y, Shou K, Zhang Q. Transplantation of the remedial free anteromedial thigh flap. *Zhonghua Zheng Xing Shao Shang Wai Ke Za Zhi* 1996;12:407-9.

10. Hayashi A, Maruyama Y. The use of the anteromedial thigh fasciocutaneous flap in the reconstruction of the lower abdomen and inguinal region: A report of two cases. *Br J Plast Surg* 1988;41:633-8.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50

**Source of Support: Nil, Conflict of Interest: None declared.**